

Liquid-gas phase behavior of polydisperse dipolar hard-sphere fluid: Extended TPT-CF.

Y. V. Kalyuzhnyi^{a)}, S. P. Hlushak^{a)}, and P. T. Cummings^{b)}

^{a)} *Institute for Condensed Matter Physics,
Sviientsitskoho 1, 79011, Lviv, Ukraine*

^{b)} *Department of Chemical Engineering, Vanderbilt University,
Nashville, Tennessee 37235,
and Nanomaterials Theory Institute,
Center for Nanophase Material Sciences, Oak Ridge National Laboratory,
Oak Ridge, Tennessee 37830*

(Dated: November 17, 2009)

Abstract

The liquid-gas phase diagram for polydisperse dipolar hard-sphere fluid with polydispersity in the hard-sphere size and dipolar moment is calculated using extension of the recently proposed thermodynamic perturbation theory for central force (TPT-CF) associating potential. To establish connection with the phase behavior of ferrocolloidal dispersions it is assumed that the dipole moment is proportional to the cube of the hard-sphere diameter. We present the full phase diagram, which includes cloud and shadow curves, binodals and distribution functions of the coexisting daughter phases at different degrees of the system polydispersity.

I. INTRODUCTION

In this paper we consider the liquid-gas phase behavior of polydisperse dipolar hard-sphere mixture. Recently, liquid-gas phase equilibria in monodisperse dipolar hard-sphere fluid, Yukawa dipolar hard-sphere fluid and Shtockmayer fluid were studied using thermodynamic perturbation theory for central force (TPT-CF) associating potential [1, 2, 3, 4]. In this study we propose extension of the TPT-CF, which enables us to investigate the phase behavior of polydisperse mixture of the dipolar hard spheres with polydispersity in both hard-sphere size and dipole moment. We call this extension as extended TPT-CF (ETPT-CF). In a certain sense ETPT-CF can be seen as a combination of Wertheim's TPT [5, 6] for associating fluid with association taking place due to off-center attractive sites and TPT-CF, which allows multiple bonding of each site. In our theory we have several Wertheim's type of associating sites with the possibility for each site to be bonded to several other sites (in Wertheim's TPT each site is only singly bondable). Final expressions for thermodynamical properties of polydisperse dipolar hard-sphere fluid is written in terms of the finite number of distribution function moments, i.e. in the framework of ETPT-CF this system belongs to the family of the so-called truncatable free energy models. This property enables us to calculate the full liquid-gas phase diagram (including cloud and shadow curves and binodals) and to study effects of fractionation on the level of the distribution functions of coexisting daughter phases.

II. EXTENDED THERMODYNAMIC PERTURBATION THEORY FOR CENTRAL FORCE ASSOCIATIVE POTENTIAL

A. Analysis and classification of diagrams

We consider a multicomponent fluid mixture consisting of n species with a number density $\rho = \sum_a^n \rho_a$ at a temperature T ($\beta = 1/k_B T$), where ρ_a is the density of the particles of a species. The particles of the species a and b interact via the pair potential $U_{ab}(12)$, which can be written as a sum of the reference $U_{ref}^{ab}(12)$ and associative $U_{ass}^{ab}(12)$ parts

$$U_{ab}(12) = U_{ref}^{ab}(12) + U_{ass}^{ab}(12), \quad (1)$$

where 1 and 2 denote positions and orientations of the particles 1 and 2. We assume that associative part of the potential can be represented as a sum of $M_a \times M_b$ terms, i.e.

$$U_{ass}^{ab}(12) = \sum_{KL} U_{KL}^{ab}(12), \quad (2)$$

where the lower indices K and L take the values $\underbrace{A, B, C, \dots}_{M_a}$ and $\underbrace{A, B, C, \dots}_{M_b}$, respectively. These values specify splitting of the total associating potential $U_{ass}^{ab}(12)$ into several particular pieces. For example in the case of the models utilized by Wertheim [6] these indices denote off-center attractive sites and in the case of the Mercedes-Benz (MB) type of the models [7] or cone models [8] they stand for the type of the hydrogen bonding arms. Hereafter we will refer to these indices as to the site indices, keeping in mind that they may have more general meaning. Here M_a and M_b are the number of such sites on the particles of a and b species, respectively. According to (1) and (2) Mayer function $f_{ab}(12)$ for the total potential (1) takes the following form:

$$f_{ab}(12) = f_{ref}^{ab}(12) + e_{ref}^{ab}(12) \left\{ \prod_{KL} [1 + f_{KL}^{ab}(12)] - 1 \right\}, \quad (3)$$

where we are using the usual notation:

$$e(12) = \exp[-\beta U(12)], \quad f(12) = e(12) - 1, \quad (4)$$

For the sake of diagrammatic analysis we will follow Wertheim [6] and instead of circles we introduce hypercircles to represent particles in diagrammatic expansions. Each hypercircle is depicted as a large open circle with small circles inside denoting the sites. Corresponding cluster integrals are represented by the diagrams build on a hypercircles connected by f_{ref} and e_{ref} bonds and a site circles connected by the associating bonds f_{KL} . Due to the decomposition of the Mayer function $f_{ab}(12)$ (3) we will have the following diagrammatic expressions for the logarithm of the grand partition function Ξ and for the one-point density $\rho_a(1)$ in terms of the activity z :

$\ln \Xi$ = sum of all topologically distinct connected diagrams consisting of field $\tilde{z}$ hypercircles, f_{ref} , e_{ref} and f_{KL} bonds. Each bonded pair of $\tilde{z}$ hypercircles has either f_{ref} , or e_{ref} and one or more f_{KL} bonds.

$\rho_a(1)$ = sum of all topologically distinct connected diagrams obtained from $\ln \Xi$ by replacing in all possible ways one field $\tilde{z}$ hypercircle by a $\tilde{z}_a(1)$ circle labeled 1.

Here $\tilde{z}_a(i) = z_a \exp[-\beta U_a(i)]$, i denotes position and orientation of the particle i , and $U_a(i)$ is an external field. For a uniform system $\tilde{z}_a(1) \equiv z_a$. Following [1, 2, 5, 6] we introduce the definition of the s -mer diagrams. These are the diagrams consisting of s hypercircles, which all are connected by the network of f_{KL} bonds. The site circles, which are incident with more than $m_K^a f_{KL}^{ab}$ bonds are called *oversaturated* site circles. We consider now the set of oversaturated site circles with each pair connected by at least one path formed by the circles from the same set. The subdiagram involving this set of circles, together with the site circles adjacent to them and f_{KL} bonds connecting all these circles, we call the *oversaturated* subdiagram. The set of all possible s -mer diagrams can be constructed in three steps: (i) generating the subset of all possible connected diagrams with only f_{KL} bonds, (ii) inserting combined bond $e_{ref} = f_{ref} + 1$ between all pairs of hypercircles with the site circles, which belong to the same maximal oversaturated subdiagram and (iii) taking all ways of inserting a f_{ref} bond between the pairs of hypercircles, which were not connected during the previous two steps. As a result the diagrams, which appear in $\ln \Xi$ and $\rho(1)$, can be expressed in terms of the s -mer diagrams:

$\ln \Xi$ = sum of all topologically distinct connected diagrams consisting of s -mer diagrams with $s = 1, \dots, \infty$ and f_{ref} bonds between pairs of hypercircles in distinct s -mer diagrams.

The procedure for obtaining the expression for $\rho_a(1)$ from $\ln \Xi$ remains unchanged.

The diagrams appearing in the $\tilde{z}$ expansion of the singlet density $\rho_a(1)$ can be classified with respect to the number of f_{KL}^{ab} bonds associated with the labeled $\tilde{z}_a(1)$

hypercircle. We denote the sum of the diagrams with $i_K \leq m_K^a$ associating bonds connected to the site K ($K = A, B, C, \dots$), which belongs to the particle of species a as $\rho_{A_{i_A}, B_{i_B}, C_{i_C}, \dots}^a(1)$. Any site K , which is connected to $i_K > m_K^a$ associating bonds, will be denoted as $K_{m_K^a}$. In what follows we will use also a condensed version of the notation, i.e.

$$\rho_{A_{i_A}, B_{i_B}, C_{i_C}, \dots}^a(1) \equiv \rho_{i_A, i_B, i_C, \dots}^a(1) \equiv \rho_{\{i\}}^a(1) \quad (5)$$

where $\{i\} = i_A, i_B, i_C, \dots$. The set $\{i\}$ with all indices, except one index i_K , equal 0, will be denoted as i_K , i.e. $\{i\} = 0, \dots, 0, i_K, 0, \dots, 0 \equiv i_K$, so that for any quantity $x_{\{i\}}^a$ we have

$$x_{\{i\}}^a = x_{0, \dots, 0, i_K, 0, \dots, 0}^a \equiv x_{K_{i_K}}^a \equiv x_{i_K}^a \quad (6)$$

Thus $\rho_a(1)$ can be written as follows

$$\rho_a(1) = \sum_{i_A, i_B, \dots=0}^{m_A^a, m_B^a, \dots} \rho_{A_{i_A}, B_{i_B}, C_{i_C}, \dots}^a(1) \equiv \sum_{\{i\}=0}^{\{m^a\}} \rho_{\{i\}}^a(1). \quad (7)$$

B. Topological reduction

The process of switching from the activity to a density expansion goes in the same fashion as in Refs. [1, 2, 6]. However, to proceed it is convenient to use an operator form of notation. The operators are introduced in a manner similar to that presented in references [1, 6] to which we refer the reader for more details. We associate with each labeled l hypercircle an operator $\epsilon_{\{i\}}^a(l)$ with the following properties:

$$\begin{aligned} \epsilon_{\{i\}}^a(l) &= 0, & \text{if any } i_K > m_K^a, \\ \epsilon_{\{i\}}^a(l) &= 1, & \text{if all } i_K = 0, \\ \epsilon_{\{i\}}^a(l) \epsilon_{\{j\}}^a(l) &= \epsilon_{\{i+j\}}^a(l), \end{aligned} \quad (8)$$

where $\{i+j\} \equiv i_A + j_A, i_B + j_B, i_C + j_C, \dots$. The one-point quantities, which, for convenience, are denoted by $x_{\{i\}}^a$, can be presented as illustrated below:

$$\hat{x}_a(1) = \sum_{\{i\}=0}^{\{m^a\}} \epsilon_{\{i\}}^a(1) x_{\{i\}}^a(1). \quad (9)$$

The operators $\epsilon_{\{i\}}^a$ are straightforward generalization of the operators introduced earlier [1, 6, 9]. Thus, the rules of manipulation with the new quantities $\hat{x}_a$ are similar to that discussed before. In particular, the usual algebraic rules apply to these quantities and analytical functions of $\hat{x}_a$ are defined by the corresponding power series. Similar, as in references [1, 6, 9], it is convenient to use the angular brackets to specify the operation

$$\langle \hat{x}_a \rangle = x_{\{m^a\}}^a. \quad (10)$$

In the case of several labeled circles the subscripts on the brackets denote the circle to which procedure (10) is to be applied.

Analyzing the connectivity of the diagrams in $\rho_a(1)$ at a labeled $\tilde{z}_a(1)$ hypercircle we have

$$\hat{\rho}_a(1)/\tilde{z}_a(1) = \exp [\hat{c}_a(1)], \quad (11)$$

where $c_{\{i\}}^a(1)$ with $\{i\} \neq \{0\}$ denotes the sum of diagrams in $\rho_{\{i\}}^a(1)/\rho_{\{0\}}^a(1)$ for which the labeled 1 hypercircle is not an articulation circle. Similarly $c_{\{0\}}^a(1)$ denotes the sum of diagrams in $\rho_{\{0\}}^a(1)/\tilde{z}_a(1)$ for which hypercircle 1 is not an articulation circle. Elimination of the diagrams containing field articulation circles can be achieved by switching from an activity to a density expansion. To do so we adopt the following rule: each field hypercircle $\tilde{z}_a$, with bonding state of its sites represented by the set $\{l\}$, in all irreducible diagrams $\hat{c}_a$ is replaced by a $\sigma_{\{j\}}^a$ hypercircle, where $j_K = m_K^a - l_K$ ($K = A, B, \dots$) for $m_K^a - l_K \geq 0$ and $j_K = 0$ for $m_K^a - l_K < 0$. The new quantities $\sigma_{\{i\}}^a(1)$ are connected to the densities $\rho_{\{i\}}^a(1)$ via the following relation:

$$\hat{\sigma}_a(1) = \hat{\rho}_a(1) \sum_{\{i\}=0}^{\{m^a\}} \epsilon_{\{i\}}^a(1). \quad (12)$$

This relation can be inverted expanding $\left[\sum_{\{i\}=0}^{\{m^a\}} \epsilon_{\{i\}}^a(1) \right]^{-1}$ into a power series, i.e.

$$\hat{\rho}_a(1) = \hat{\sigma}_a(1) \prod_{K=A} [1 - \epsilon_{K_1}^a(1)] \quad (13)$$

Now the diagrammatic expansions for $c_{\{i\}}^a$ can be expressed in terms of irreducible diagrams. To present this result in compact and convenient form we introduce a sum of the diagrams $c^{(0)}$ defined as follows:

$c^{(0)}$ = sum of all topologically distinct irreducible diagrams consisting of s -mer diagrams with $s = 1, \dots, \infty$ and f_{ref} bonds between pairs of hypercircles in distinct s -mer diagrams. All hypercircles are field circles carrying the σ -factor according to the rule formulated above

Functional differentiation of $c^{(0)}$ with respect to $\sigma_{\{m^a-i\}}^a$ gives an expression for $c_{\{i\}}^a$:

$$c_{\{i\}}^a = \frac{\delta c^{(0)}}{\delta \sigma_{\{m^a-i\}}^a}. \quad (14)$$

C. Extended thermodynamic perturbation theory for central force associating potential

Now we are in a position to rewrite the regular one-density virial expansion for the pressure P in terms of the density parameters $\hat{\sigma}_a(1)$. Following the scheme, proposed earlier [2, 4, 5, 6] we have expression for the pressure in operator form

$$\beta PV = \sum_a \int \langle \hat{\sigma}_a(1) [1 - \hat{c}_a(1)] \rangle d(1) + c^{(0)} \quad (15)$$

and explicitly

$$\beta PV = \sum_a \int \left[\rho_a(1) - \sum_{\{i\}=0}^{\{m^a\}} \sigma_{\{m^a-i\}}^a(1) c_{\{i\}}^a(1) \right] d(1) + c^{(0)}, \quad (16)$$

where V is the volume of the system. Similarly, as in [5, 6] one can verify that these expressions satisfy the regular thermodynamic relation $\bar{\rho}_a = \beta \partial P / \partial \mu_a$, where $\bar{\rho}_a = \int \rho_a(1) d(1)$ and μ^a is the chemical potential. This can be achieved by taking a variation of (15) (or (16)) and combining (11), (13) and (14). The corresponding expression for Helmholtz free energy is

$$\beta A = \sum_a \int \left[\rho_a(1) \ln \frac{\rho_{\{0\}}^a(1)}{\Lambda_a} + \sum_{\{i\} \neq 0}^{\{m^a\}} \sigma_{\{m^a-i\}}^a(1) c_{\{i\}}^a(1) \right] d(1) - c^{(0)}, \quad (17)$$

where Λ_a is the thermal de Broglie wavelength. This expression is derived using the regular thermodynamic expression for Helmholtz free energy $A = \sum_a N_a \mu_a - PV$ together with relation

$$\beta N_a \mu_a = \int \rho_a(1) \left[\ln \frac{\rho_{\{0\}}^a(1)}{\Lambda_a} - c_{\{0\}}^a(1) \right] d(1), \quad (18)$$

which follows from (11), written for $\rho_{\{0\}}^a$. Here N_a is the number of particles of species a in the system.

Helmholtz free energy in excess to its reference system value A_{ref} is obtained by subtracting corresponding expression for A_{ref} from (17), i.e.

$$\begin{aligned} & \beta (A - A_{ref}) = \\ & = \sum_a \int \left[\rho_a(1) \ln \frac{\rho_{\{0\}}^a(1)}{\rho_a(1)} + \sum_{\{i\} \neq 0}^{\{m^a\}} \sigma_{\{m^a-i\}}^a(1) c_{\{i\}}^a(1) \right] d(1) - \left(c^{(0)} - c_{ref}^{(0)} \right), \end{aligned} \quad (19)$$

where $c_{ref}^{(0)}$ is the corresponding sum of the diagrams for the reference system. Ordering the virial expansion (19) with respect to the number of associating f_{KL} bonds and neglecting the terms with more than one associating bond we have

$$c^{(0)} - c_{ref}^{(0)} = \frac{1}{2} \sum_{ab} \int g_{ref}^{ab}(12) \langle \hat{\sigma}_a(1) \hat{f}_{ab}(12) \hat{\sigma}_b(2) \rangle_{12} d(1) d(2) \quad (20)$$

and

$$\hat{c}_a(1) - c_{\{0\}}^a(1) = \sum_b \int g_{ref}^{ab}(12) \langle \hat{f}_{ab}(12) \hat{\sigma}_a(2) \rangle_2 d(2), \quad (21)$$

where $g_{ref}^{ab}(12)$ is the reference system distribution function and

$$\hat{f}_{ab}(12) = \sum_{KL} \epsilon_{K_1}^a(1) f_{KL}^{ab}(12) \epsilon_{L_1}^b(2). \quad (22)$$

Due to single bond approximation $c_{\{i\}}^a = 0$ for all values of the set $\{i\}$, except for $\{i\} = 0$ and $\{i\} = i_K$ with $i_K = 1$. This property together with (11) yield the following relations:

$$c_{K_1}^a(1) = \rho_{K_1}^a(1) / \rho_{\{0\}}^a(1) \quad (23)$$

and

$$\frac{\rho_{\{i\}}^a(1)}{\rho_{\{0\}}^a(1)} = \prod_K \frac{1}{i_K!} [c_{K_1}^a(1)]^{i_K} = \prod_K \frac{1}{i_K!} \left[\frac{\rho_{K_1}^a(1)}{\rho_{\{0\}}^a(1)} \right]^{i_K}, \quad \text{for } i_K \in \{i\} \quad (24)$$

The set of relations (20), (21) and (23) defined all the quantities needed to calculate the Helmholtz free energy of the system (19), provided that the properties of the reference system are known.

Finally it is worth noting, that ETPT-CF theory developed here reduces to TPT1 proposed by Wertheim [6], if for all sites single-bonding condition $m_K^a = 1$ is assumed. In the other limiting case of only one site per particle ETPT-CF will coincide with TPT-CF developed earlier [1, 2, 4].

D. Extended TPT-CF for two sites with double-bonding condition

The theory presented in the previous section is quite general and can be applied to a number of different situations. However in the present study we are interested in the version of the theory for the model with two sites both of which can be bonded twice. More specifically we are interested in the extension and application of the theory to study the phase behavior of polydisperse dipolar hard-sphere mixture.

We assume that each of the particles in the system have two doubly bondable attractive sites, A and B , i.e. we have: $M_a = 2$ and $m_A^a = m_B^a$. We also assume, that attractive interaction is acting only between the sites of the same sort. Using these suggestions, relations (11) and (12) and taking into account that the system is uniform, the density parameters $\sigma_{A_2B_2}^a = \rho_a \sigma_{A_1B_2}^a \equiv \tilde{\sigma}_{A_1}^a$ and $\sigma_{A_2B_1}^a \equiv \tilde{\sigma}_{B_1}^a a$ can be expressed in terms of $c_{K_1}^a$

$$\rho_a = \frac{1}{4} \sigma_{A_0B_0}^a \left[1 + (\kappa_{A_1}^a)^2 \right] \left[1 + (\kappa_{B_1}^a)^2 \right], \quad (25)$$

$$\tilde{\sigma}_{A_1}^a = \frac{1}{2} \sigma_{A_0B_0}^a \kappa_{A_1}^a \left[1 + (\kappa_{B_1}^a)^2 \right], \quad (26)$$

$$\tilde{\sigma}_{B_1}^a = \frac{1}{2} \sigma_{A_0B_0}^a \kappa_{B_1}^a \left[1 + (\kappa_{A_1}^a)^2 \right], \quad (27)$$

where K takes the values A and B and $\kappa_{K_1}^a = 1 + c_{K_1}^a$. These two equations give

$$\sigma_{A_0B_0}^a = 4\rho_a \left\{ \left[1 + (\kappa_{A_1}^a)^2 \right] \left[1 + (\kappa_{B_1}^a)^2 \right] \right\}^{-1}, \quad (28)$$

and

$$\tilde{\sigma}_{K_1}^a = \frac{2\rho_a \kappa_{K_1}^a}{1 + (\kappa_{K_1}^a)^2}. \quad (29)$$

In turn, using (21), for $\kappa_{K_1}^a$ we have

$$\kappa_{K_1}^a = 1 + \sum_b I_{KK}^{ab} \tilde{\sigma}_{K_1}^a, \quad (30)$$

where

$$I_{KK}^{ab} = \int g_{ref}^{ab}(12) f_{KK}^{ab}(12) d(2). \quad (31)$$

Combining (20), (29) and (30) expression for Helmholtz free energy (19) can be written in terms of $\kappa_{K_1}^a$ parameters

$$\beta \frac{A - A_{ref}}{V} = \sum_a \rho_a \left[\ln \frac{\rho_{A_0 B_0}^a}{\rho_a} - \kappa_{A_1}^a \frac{1 - \kappa_{A_1}^a}{1 + (\kappa_{A_1}^a)^2} - \kappa_{B_1}^a \frac{1 - \kappa_{B_1}^a}{1 + (\kappa_{B_1}^a)^2} \right], \quad (32)$$

which satisfy the following set of equations:

$$\kappa_{K_1}^a = 1 + \sum_b \frac{2\rho_b \kappa_{K_1}^b}{1 + (\kappa_{K_1}^b)^2} I_{KK}^{ab} \quad (33)$$

Chemical potential $\Delta\mu_a$ and pressure ΔP in excess to their reference system values can be obtained using standard thermodynamical relations:

$$\mu_a - \mu_{ref}^a = \frac{\partial [(A - A_{ref})/V]}{\partial \rho_a}, \quad P - P_{ref} = \sum_a \rho_a (\mu_a - \mu_{ref}^a) - \frac{A - A_{ref}}{V}. \quad (34)$$

Finally, the average size of the clusters, which appear in the system, can be characterized by the average length of the chain L_K formed by either A -bonded ($K = A$) or B -bonded ($K = B$) particles. This quantity is defined by the following expression

$$L_K = \frac{\sum_a (\alpha_{K,end}^a + \alpha_{K,mid}^a + \alpha_0^a)}{\sum_a (\alpha_{K,end}^a/2 + \alpha_0^a)}, \quad (35)$$

where $\alpha_{K,end}^a$ is the fraction of singly K -bonded particles (fraction of the chain ends), $\alpha_{K,mid}^a$ is the fraction of doubly K -bonded particles (fraction of the chain middles) and α_0^a is the fraction of nonbonded particles. For these fractions we have

$$\rho \alpha_{A,end}^a = \tilde{\sigma}_{A_1}^a - \sigma_{A_0 B_2}^a, \quad \rho \alpha_{A,mid}^a = \rho - \tilde{\sigma}_{A_1}^a, \quad \rho \alpha_0^a = \sigma_{A_0 B_0}^a. \quad (36)$$

Substituting these expressions into expression for L_A (35) and using (28), (29) and expression for $\sigma_{A_0 B_2}^a$,

$$\sigma_{A_0 B_2}^a = \frac{1}{2} \rho_a \sigma_{A_0 B_0}^a \left[(\kappa_{B_1}^a)^2 + 1 \right], \quad (37)$$

we get the final expression for L_A in terms of $\kappa_{K_1}^a$:

$$L_K = \sum_a \rho_a \frac{4 - [1 + (\kappa_{K_1}^a)^2] [1 - (\kappa_{K_1}^a)^2]}{[1 + (\kappa_{A_1}^a)^2] [1 + (\kappa_{B_1}^a)^2]} \left\{ \sum_a \rho_a \frac{4 - (1 - \kappa_{K_1}^a) [1 + (\kappa_{K_1}^a)^2]}{[1 + (\kappa_{A_1}^a)^2] [1 + (\kappa_{B_1}^a)^2]} \right\}^{-1}, \quad (38)$$

where if $K = A$ then $\tilde{K} = B$ and if $K = B$ then $\tilde{K} = A$.

III. LIQUID-GAS PHASE BEHAVIOR OF POLYDISPERSE DIPOLAR HARD-SPHERE FLUID

A. The model

We consider polydisperse dipolar hard-sphere fluid mixture with a number density ρ and polydispersity in both hard-sphere diameter σ and dipolar moment d_μ . We assume, that the dipole moment is proportional to the particle volume, i.e. $d_\mu \sim \sigma^3$. Thus the type of the particle is completely defined by its hard-sphere size and hereafter we will be using σ instead of the indices $a, b, \dots$ to denote the particle species. We also assume, that hard-sphere size of the particles is distributed according to normalized distribution function $F(\sigma) \geq 0$,

$$\int_0^\infty F(\sigma) d\sigma = 1. \quad (39)$$

Interaction between particles of species σ_1 and σ_2 in our system is described by the following pair potential:

$$U(r, \sigma_1 \sigma_2) = U_{hs}(r, \sigma_1 \sigma_2) + U_{dd}(12, \sigma_1 \sigma_2), \quad (40)$$

where $U_{hs}(r, \sigma_1 \sigma_2)$ is the hard-sphere potential and $U_{dd}(r, \sigma_1 \sigma_2)$ is the dipole-dipole potential, given by

$$U_{dd}(12, \sigma_1 \sigma_2) = -\frac{d_\mu(\sigma_1)d_\mu(\sigma_2)}{r^3} [2 \cos \varphi_1 \cos \varphi_2 - \sin \varphi_1 \sin \varphi_2 \cos(\phi_1 - \phi_2)]. \quad (41)$$

Here φ_1 and φ_2 denote the angles between the dipole vectors and vector, which joins the centers of the two particles, and ϕ_1 and ϕ_2 are the azimuthal angles about this vector. To proceed we have to split the total potential (40) into the reference and associative pieces. We assume, that the reference part of the potential is represented by the hard-sphere part $U_{hs}(r, \sigma_1 \sigma_2)$ and associative part by the dipole-dipole potential $U_{dd}(r, \sigma_1 \sigma_2)$. At the contact distance $\sigma_{12} = (\sigma_1 + \sigma_2)/2$ the latter potential has two equal potential minima of the depth $-2d_\mu(\sigma_1)d_\mu(\sigma_2)/\sigma_{12}^3$ at “nose-to-tail” configuration ($\varphi_1 = \varphi_2 = 0$, $\varphi_1 = \varphi_2 = \pi$). These minima are responsible for the formation of the chains of the particles in the system. In addition there

are twice less deep minima $(-d_\mu(\sigma_1)d_\mu(\sigma_2)/\sigma_{12}^3)$ at antiparallel configuration with $\varphi_1 = \varphi_2 = \pi/2$, $\phi_1 - \phi_2 = \pi$. The latter minima cause the formation of the network connecting the chains. According to earlier theoretical and computer simulation studies [10, 11] competition between chain formation and network formation defines the existence of the liquid-gas phase transition in the dipolar hard-sphere fluid. To account for this effect we propose the following splitting of the total associative potential $U_{ass}(12, \sigma_1\sigma_2) = U_{dd}(12, \sigma_1\sigma_2)$:

$$U_{BB}(12, \sigma_1\sigma_2) = \Theta(\varphi_1) \Theta(\varphi_2) U_{dd}(12, \sigma_1\sigma_2), \quad (42)$$

$$U_{AA}(12, \sigma_1\sigma_2) = [1 - \Theta(\varphi_1) \Theta(\varphi_2)] U_{dd}(12, \sigma_1\sigma_2), \quad (43)$$

where $\Theta(\varphi) = H(\pi/2 + \varphi_0 - \varphi) H(\pi/2 - \varphi_0 + \varphi)$ and $H(x)$ is the Heaviside step function. Here φ_0 plays a role of the potential splitting parameter. For $\varphi_0 = \pi/2$ we have that $U_{BB}(12, \sigma_1\sigma_2) = U_{dd}(12, \sigma_1\sigma_2)$ and $U_{AA}(12, \sigma_1\sigma_2) = 0$. On the other hand $\varphi_0 = 0$ gives: $U_B(12, \sigma_1\sigma_2) = 0$ and $U_{AA}(12, \sigma_1\sigma_2) = U_{dd}(12, \sigma_1\sigma_2)$. In both limiting cases the theory developed will treat the system as a polydisperse mixture of the hard-sphere chains. For the intermediate values of φ_0 the energy minima at “nose-to-tail” configuration are included into $U_{AA}(12, \sigma_1\sigma_2)$ and network forming minima appear in $U_{BB}(12, \sigma_1\sigma_2)$. Our choice for the particular value of φ_0 will be discussed later.

B. Thermodynamic properties

For a general multicomponent dipolar hard-sphere mixture thermodynamic properties can be obtained using solution of the set of nonlinear equations (33) and expression for Helmholtz free energy (32). However even for multicomponent case solution of this equation rapidly becomes involved as the number of components increases. As we proceed to polydisperse case solution of the polydisperse version of equation (33) becomes intractable, since now we have to deal with the following integral equation:

$$\kappa_K(\sigma_1) = 1 + 2\rho \int_0^\infty F(\sigma_2) \frac{\kappa_K(\sigma_2) I_{KK}(\sigma_1\sigma_2)}{1 + \kappa_K^2(\sigma_2)} d\sigma_2, \quad (44)$$

where we have dropped the lower index 1, i.e. $\kappa_{K1}(\sigma) \equiv \kappa_K(\sigma)$. In order to solve this equation we propose here to interpolate the key quantity of the theory, the

volume integral $I_{KL}(\sigma_1\sigma_2)$, using a sum of N_Y Yukawa terms. Since the reference system pair distribution function $g_{ref}(r, \sigma_1\sigma_2)$ is independent of mutual orientation of the particles for the integral (31) we have

$$I_{KK}(\sigma_1\sigma_2) = 4\pi \int_0^\infty r^2 g_{ref}(r, \sigma_1\sigma_2) \bar{f}_{KK}(r, \sigma_1\sigma_2) dr, \quad (45)$$

where $\bar{f}_{KK}(r, \sigma_1\sigma_2)$ is orientation averaged Mayer function for associative potential $U_{KK}(12, \sigma\sigma)$. We assume, that $\bar{f}_{KK}(r, \sigma_1\sigma_2)$ can be represented in the following form:

$$\bar{f}_{KK}(r, \sigma_1\sigma_2) = \frac{1}{4\pi r} \sum_n^{N_Y} A_K^{(n)}(\sigma_1) A_K^{(n)}(\sigma_2) e^{-z_K^{(n)}(r-\sigma_{12})}. \quad (46)$$

Parameters $A_K^{(n)}(\sigma)$ and $z_K^{(n)}$ are obtained using the interpolation scheme, which is presented and discussed in the Appendix A. Using (45) and (46), we have

$$I_{KK}(\sigma_1\sigma_2) = \sum_n^{N_Y} A_K^{(n)}(\sigma_1) A_K^{(n)}(\sigma_2) G_K^{(n)}(\sigma_1\sigma_2), \quad (47)$$

where $G_K^{(n)}(\sigma_1\sigma_2) = \int_0^\infty r e^{-z_K^{(n)}r} g_{ref}(r, \sigma_1\sigma_2) dr$ is the Laplace transform of the radial distribution function $g_{ref}(r, \sigma_1\sigma_2)$. We will be using here Percus-Yewick approximation for hard-sphere radial distribution function, since analytical expression for its Laplace transform is known [12]

$$e^{-z_K^{(n)}\sigma_{12}} G_K^{(n)}(\sigma_1\sigma_2) = \frac{\Delta}{(z_K^{(n)})^2 D_K^{(n)}} \left[z_K^{(n)} \left(\sigma_{12} + \sigma_1\sigma_2 \frac{\pi}{4\Delta} m_2 \right) + 1 + \frac{\pi}{2\Delta} m_3 + \right. \\ \left. + \frac{\pi z_K^{(n)}}{2\Delta} \left(m_{K,2}^{(n)} - 2\sigma_{12} m_{K,1}^{(n)} + \sigma_1\sigma_2 m_{K,0}^{(n)} \right) \right], \quad (48)$$

where

$$D_K^{(n)} = \Delta^2 - \frac{2\pi}{z_K^{(n)}} \left(1 + \frac{1}{2}\pi m_3 \right) \left(m_{K,0}^{(n)} + \frac{1}{2}m_2 \right) - \\ - 2\pi \left\{ \Delta m_{K,1}^{(n)} + \frac{1}{4}\pi \left[m_{K,2}^{(n)} \left(m_2 + 2m_{K,0}^{(n)} \right) - \left(m_{K,1}^{(n)} \right)^2 \right] \right\}, \quad (49)$$

$$\Delta = 1 - \pi m_3/6. \quad (50)$$

Here m_l are the moments and $m_{K,l}^{(n)}$ are the generalized moments of the distribution function $F(\sigma)$. Symbolically expression for these moments can be represented as follows:

$$m = \int_0^\infty m(\sigma) F(\sigma) d\sigma. \quad (51)$$

Hereafter all the quantities denoted as m with certain set of indices will represent the generalized moments defined by (51). Corresponding expressions for $m(\sigma)$ are collected in the Appendix B. Inserting (47) into equation (44) and using (48), we find

$$\kappa_K(\sigma) = 1 + \rho \sum_n \sum_{j=1}^{N_Y} C_{K,j}^{(n)} \Omega_{K,j}^{(n)}(\sigma), \quad (52)$$

where $C_{K,j}^{(n)}$ satisfies the following set of equations:

$$C_{K,j}^{(n)} = 2 \int_0^\infty \sigma^{j-1} F(\sigma) A_K^{(n)}(\sigma) \frac{1 + \rho \sum_l \sum_{i=1}^2 \Omega_{K,i}^{(l)}(\sigma) C_{K,i}^{(l)}}{1 + \left[1 + \rho \sum_l \sum_{i=1}^2 \Omega_{K,i}^{(l)}(\sigma) C_{K,i}^{(l)} \right]^2} d\sigma. \quad (53)$$

Here

$$\Omega_{K,1}^{(n)}(\sigma) = A_K^{(n)}(\sigma) \left(\sigma P_{K,1}^{(n)} + P_{K,3}^{(n)} \right), \quad \Omega_{K,2}^{(n)}(\sigma) = A_K^{(n)}(\sigma) \left(\sigma P_{K,2}^{(n)} + P_{K,1}^{(n)} \right), \quad (54)$$

$$P_{K,1}^{(n)} = \frac{\Delta}{2z_K^{(n)} D_K^{(n)}} \left(1 - \frac{\pi}{\Delta} m_{K,1}^{(n)} \right), \quad (55)$$

$$P_{K,2}^{(n)} = \frac{\pi}{2z_K^{(n)} D_K^{(n)}} \left(\frac{1}{2} m_2 + m_{K,0}^{(n)} \right), \quad (56)$$

$$P_{K,3}^{(n)} = \frac{\Delta}{\left(z_K^{(n)} \right)^2 D_K^{(n)}} \left[1 + \frac{\pi}{2\Delta} \left(m_3 + z_K^{(n)} m_{K,2}^{(n)} \right) \right], \quad (57)$$

Thus solution of the integral equation (44) for the unknown function $\kappa_K(\sigma)$ now is reduced to solution of a set of equations for $4N_Y$ unknown constants $C_{K,j}^{(n)}$. This solution can be used to calculate $\kappa_K^{(n)}(\sigma)$, which in turn can be utilized to calculate thermodynamical properties of the system via Helmholtz free energy (32). Generalizing expression for Helmholtz free energy (32) for polydisperse system, we have

$$\beta \frac{A - A_{ref}}{V} = \rho \int_0^\infty F(\sigma) \left[-\ln \frac{1}{4} \prod_{K=A}^B (\kappa_A^2(\sigma) + 1) - \sum_{K=A}^B \kappa_K(\sigma) \frac{1 - \kappa_K(\sigma)}{1 + \kappa_K^2(\sigma)} \right] d\sigma. \quad (58)$$

Now we can use the standard relation between Helmholtz free energy and chemical potential (34), generalized to polydisperse case

$$\beta [\mu(\sigma) - \mu_{ref}(\sigma)] = \frac{\beta}{\rho} \frac{\delta \{A - A_{ref}/V\}}{\delta \{F(\sigma)\}}, \quad (59)$$

where $\delta/\delta \{F(\sigma)\}$ denote functional differentiation with respect to the distribution $F(\sigma)$. We find

$$\beta [\mu(\sigma) - \mu_{ref}(\sigma)] = \frac{m_\mu}{\rho} - \ln \frac{1}{4} \prod_{K=A}^B (\kappa_A^2(\sigma) + 1) - \sum_{K=A}^B \kappa_K(\sigma) \frac{1 - \kappa_K(\sigma)}{1 + \kappa_K^2(\sigma)}$$

$$+ \sum_n^{N_Y} \sum_{K=A}^B \left(\sum_{j=1}^3 \mu_{K,j}^{(n)} \frac{\delta P_{K,j}^{(n)}}{F(\sigma)} + \sum_{j=1}^2 \nu_{K,j}^{(n)} \frac{\delta C_{K,j}^{(n)}}{F(\sigma)} \right). \quad (60)$$

Here

$$\mu_{K,1}^{(n)} = \frac{1}{2} \left(m_{K,0}^{(n,0)} C_{K,2}^{(n)} + m_{K,1}^{(n,0)} C_{K,1}^{(n)} \right), \quad (61)$$

$$\mu_{K,2}^{(n)} = \frac{1}{2} \left(m_{K,1}^{(n,0)} - \rho C_{K,2}^{(n)} \right) C_{K,2}^{(n)}, \quad \mu_{K,3}^{(n)} = \frac{1}{2} \left(m_{K,0}^{(n,0)} - \rho C_{K,1}^{(n)} \right) C_{K,1}^{(n)}, \quad (62)$$

$$\nu_{K,1}^{(n)} = \frac{1}{2} \left[\left(m_{K,1}^{(n,0)} - \rho C_{K,2}^{(n)} \right) P_{K,1}^{(n)} + \left(m_{K,0}^{(n,0)} - \rho C_{K,1}^{(n)} \right) P_{K,3}^{(n)} \right], \quad (63)$$

$$\nu_{K,2}^{(n)} = \frac{1}{2} \left[\left(m_{K,0}^{(n,0)} - \rho C_{K,1}^{(n)} \right) P_{K,1}^{(n)} + \left(m_{K,1}^{(n,0)} - \rho C_{K,2}^{(n)} \right) P_{K,2}^{(n)} \right], \quad (64)$$

functional derivatives $\delta P_{K,j}^{(n)}/\delta F(\sigma)$ and $\delta D_K^{(n)}/\delta F(\sigma)$ are presented in the Appendix B and functional derivatives $\delta C_{K,j}^{(n)}/\delta F(\sigma)$ can be obtained from the solution of the set of linear equations, which follows from (53) upon its functional differentiation with respect to $F(\sigma)$, i.e.

$$\sum_{j=1}^2 \mathbf{M}_{i,j}^{(K)} \frac{\delta \mathbf{C}_j^{(K)}}{\delta F(\sigma)} = \mathbf{R}_i^{(K)}(\sigma), \quad i = 1, 2, \quad K = A, B, \quad (65)$$

where $\left[\delta \mathbf{C}_j^{(K)}/\delta F(\sigma) \right]_n \equiv \delta C_{K,j}^{(n)}/\delta F(\sigma)$ and the elements of the matrices $\mathbf{M}_{i,j}^{(K)}$ and $\mathbf{R}_i^{(K)}(\sigma)$ are collected in the Appendix B.

The pressure expression follows from (34), generalized to polydisperse case

$$P - P_{ref} = \rho \int_0^\infty F(\sigma) [\mu(\sigma) - \mu_{ref}(\sigma)] d\sigma - \frac{A - A_{ref}}{V}. \quad (66)$$

Using this expression together with expression for the chemical potential (60), we find

$$\beta (P - P_{ref}) = m_\mu + \rho \sum_n^{N_Y} \sum_{K=A}^B \left(\sum_{j=1}^3 \mu_{K,j}^{(n)} J_{K,j}^{(n)} + \sum_{j=1}^2 \nu_{K,j}^{(n)} S_{K,j}^{(n)} \right), \quad (67)$$

where

$$J_{K,j}^{(n)} = \int_0^\infty F(\sigma) \frac{\delta P_{K,j}^{(n)}}{\delta F(\sigma)} d\sigma, \quad S_{K,j}^{(n)} = \int_0^\infty F(\sigma) \frac{\delta C_{K,j}^{(n)}}{\delta F(\sigma)} d\sigma. \quad (68)$$

Expression for the integral $J_{K,j}^{(n)}$ is presented in the Appendix B and integral $S_{K,j}^{(n)}$ can be obtain from the solution of the set of linear equations

$$\sum_{j=1}^2 \mathbf{M}_{i,j}^{(K)} \mathbf{S}_j^{(K)} = \mathbf{E}_i^{(K)}, \quad i = 1, 2, \quad K = A, B, \quad (69)$$

which follows from the set of equations (65). Here $\left[\mathbf{S}_j^{(K)} \right]_n \equiv S_{K,j}^{(n)}$ and the elements of the matrix $\mathbf{E}_i^{(K)}$ are presented in the Appendix B.

Expressions for the chemical potential (60) and pressure (67) are the final expressions to be used in the phase equilibrium calculations. The properties of the reference system (chemical potential $\mu_{ref}(\sigma)$ and pressure P_{ref}) are described here using polydisperse versions of the Mansoori *et al.* [13] expressions

$$\beta\mu_{ref}^{(ex)}(\sigma) = \left[\left(\sigma \frac{m_2}{m_3} \right)^2 \left(3 - 2\sigma \frac{m_2}{m_3} \right) - 1 \right] \ln \Delta + m_2 \left(1 + \frac{m_2\sigma}{m_3\Delta} \right) + \frac{\pi\sigma}{2\Delta} \left\{ \frac{1}{3}\sigma^2 \left[\rho - \frac{m_2^3}{m_3^2} \frac{1+\Delta}{\Delta} + \frac{\pi}{\Delta} m_2 \left(\frac{1}{2}m_1 + \frac{1}{3} \frac{m_2^2}{m_3\Delta} \right) \right] + \sigma m_1 \right\}, \quad (70)$$

$$\beta P_{ref} = \frac{1}{\Delta} \left[\rho + \frac{\pi}{2\Delta} m_1 m_2 + \frac{\pi^2}{12\Delta^2} m_2^3 - \left(\frac{\pi}{6} \right)^3 \frac{1}{\Delta^2} m_3 m_2^3 \right], \quad (71)$$

where $\mu_{ref}^{(ex)}(\sigma)$ is the reference system chemical potential in excess to its ideal gas value.

C. Phase equilibrium conditions

One can easily see that the thermodynamical properties of the model at hand obtained above are defined by the set of the finite number of the distribution function moments, i.e., four regular moments (m_l , $l = 0, 1, 2, 3$) and $1 + 10N_Y + 3N_Y(N_Y + 1)$ generalized moments (m_μ , $m_{K,i}^{(n)}$, $m_{K,j}^{(n0)}$, $m_{K,l}^{(nm)}$; $i = 0, 1, 2$; $j = 0, 1$; $l = 0, 1, 2$; $K = A, B$). Note, that $m_{K,l}^{(nm)} = m_{K,l}^{(mn)}$. Thus the polydisperse mixture of dipolar hard spheres treated within ETPT-CF belongs to the class of truncatable free energy models [14]. This property allows us to map the phase coexistence relations onto a set on nonlinear equations for the unknown moments of the daughter distribution functions.

We assume that at a certain density ρ_0 and composition $F_0(\sigma)$ the system separates into two phases with the densities ρ_1 and ρ_2 , and compositions $F_1(\sigma)$ and $F_2(\sigma)$. Hereafter the lower index 0 refer to the parent phase and the lower indices 1 and 2 refer to the daughter phases. At equilibrium these quantities take the values, which follows from the phase equilibrium conditions, i.e.: (i) conservation of the total volume of the system, (ii) conservation of the total number of the particles of each species, (iii) equality of the chemical potentials of particles of the same species in the coexisting phases, (iv) equality of the pressure in the coexisting phases. These

conditions finally lead to the following set of relations [15, 16]:

$$F_\alpha(\sigma) = F_0(\sigma) Q_\alpha(\sigma; \rho_0, \rho_1, \rho_2; [F_\alpha]), \quad (72)$$

$$P_1(\rho_1; [F_1]) = P_2(\rho_2; [F_2]), \quad (73)$$

$$\int_0^\infty F_\alpha(\sigma) d\sigma = 1, \quad \text{for } \alpha = 1 \text{ or } \alpha = 2, \quad (74)$$

where

$$\rho_\alpha Q_\alpha(\sigma; \rho_0, \rho_1, \rho_2; [F_\alpha]) = \frac{\rho_0(\rho_2 - \rho_1) \{1 - \delta_{1\alpha} + \delta_{1\alpha} \exp(\beta \Delta\mu^{(ex)})\}}{\rho_0 - \rho_1 - (\rho_0 - \rho_2) \exp(\beta \Delta\mu^{(ex)})}, \quad (75)$$

$$\Delta\mu^{(ex)} = \mu_2^{(ex)}(\sigma, \rho_2; [F_2]) - \mu_1^{(ex)}(\sigma, \rho_1; [F_1]), \quad (76)$$

$\mu_\alpha^{(ex)}$ is the excess (over the ideal gas) chemical potential of the particle of species σ in the phase α and $[\dots]$ denote functional dependence. The relation between $F_0(\sigma)$ and daughter phase distribution function $F_\alpha(\sigma)$, i.e., Eq. (72), follows from the phase equilibrium conditions (i)-(iii).

Relations (72)-(74) represent a closed set of equations to be solved for the unknowns ρ_α and $F_\alpha(\sigma)$; this set have to be solved for every value of the species variable σ . However, since thermodynamical properties of the model at hand are defined by the finite number of the moments we can map this set of equations onto a closed set of $10 + 28N_Y + 6N_Y(N_Y + 1)$ algebraic equations for ρ_α , $C_{K,1}^{(n)(\alpha)}$, $C_{k,2}^{(n)(\alpha)}$ and moments $m_k^{(\alpha)}$, $m_\mu^{(\alpha)}$, $m_{K,i}^{(n)(\alpha)}$, $m_{K,j}^{(n0)(\alpha)}$, $m_{K,l}^{(nm)(\alpha)}$ in the two coexisting phases ($\alpha = 1, 2$). We have

$$m_k^{(\alpha)} = \rho_\alpha \int_0^\infty m_k(\sigma) F_0(\sigma) Q_\alpha(\sigma, \rho_0; \{X_1\}, \{X_2\}) d\sigma, \quad k = 0, 1, 2, 3, \quad (77)$$

$$m_\mu^{(\alpha)} = \rho_\alpha \int_0^\infty m_\mu^{(\alpha)}(\sigma) F_0(\sigma) Q_\alpha(\sigma, \rho_0; \{X_1\}, \{X_2\}) d\sigma, \quad (78)$$

$$m_{K,i}^{(n)(\alpha)} = \rho_\alpha \int_0^\infty m_{K,i}^{(n)(\alpha)}(\sigma) F_0(\sigma) Q_\alpha(\sigma, \rho_0; \{X_1\}, \{X_2\}) d\sigma, \quad i = 0, 1, 2, \quad (79)$$

$$m_{K,j}^{(n0)(\alpha)} = \rho_\alpha \int_0^\infty m_{K,j}^{(n0)(\alpha)}(\sigma) F_0(\sigma) Q_\alpha(\sigma, \rho_0; \{X_1\}, \{X_2\}) d\sigma, \quad j = 0, 1, \quad (80)$$

$$m_{K,l}^{(nm)(\alpha)} = \rho_\alpha \int_0^\infty m_{K,l}^{(nm)(\alpha)}(\sigma) F_0(\sigma) Q_\alpha(\sigma, \rho_0; \{X_1\}, \{X_2\}) d\sigma, \quad l = 0, 1, 2, \quad (81)$$

where $K = A, B$ and $\{X_\alpha\}$ represent unknowns of the problem, i.e.

$$\{X_\alpha\} = \left\{ \rho_\alpha, m_k^{(\alpha)}, m_\mu^{(\alpha)}, m_{K,i}^{(n)(\alpha)}, m_{K,j}^{(n0)(\alpha)}, m_{K,l}^{(nm)(\alpha)} \right\}, \quad \alpha = 1, 2.$$

The remaining $2 + 8N_Y$ equations follows from the equality of the pressure in coexisting phases (73),

$$P_1(\rho_1; \{X_1\}) = P_2(\rho_2; \{X_2\}), \quad (82)$$

from the set of equations for $C_{K,1}^{(n)(\alpha)}$ and $C_{K,2}^{(n)(\alpha)}$ (53) and from the normalizing condition (74) for either phase $\alpha = 1$ or $\alpha = 2$,

$$\int_0^\infty F_0(\sigma) Q_\alpha(\sigma, \rho_0; \{X_1\}, \{X_2\}) d\sigma = 1. \quad (83)$$

Solution of the set of equations (53), (77)-(83) for a given density ρ_0 and distribution function $F_0(\sigma)$ of the parent phase gives the densities ρ_α and distribution functions $F_\alpha(\sigma)$ of the two coexisting daughter phases. The coexisting densities at different densities of the parent phase ρ_0 defined binodals, which are terminated when a density of one of the phases is equal to the parent phase density ρ_0 . These termination points form the cloud and shadow coexisting curves. These curves intersect at the critical point, which is characterized by the critical density $\rho_{cr} = \rho_1 = \rho_2 = \rho_0$ and critical temperature T_{cr} . The cloud-shadow curves can be obtained as a special solution of the general coexisting problem, when the properties of one phase are equal to the properties of the parent phase: assuming that the phase $\alpha = 2$ is the cloud phase, i.e. $\rho_2 = \rho_0$, and following the above scheme we will end up with the same set of equations (53), (77)-(83), but with ρ_2 and $F_2(\sigma)$ substituted by ρ_0 and $F_0(\sigma)$, respectively.

IV. RESULTS AND DISCUSSION

In this section we present our numerical results for the liquid-gas phase diagram of polydisperse dipolar hard-sphere fluid at different degree of polydispersity. For size distribution function $F(\sigma)$ we have chosen the beta distribution given by

$$F(\sigma) = \frac{\Gamma(\nu + \mu - 1)}{\Gamma(\nu)\Gamma(\mu)} \frac{(1-x)^{\mu-1} x^{\nu-1}}{(\sigma_u - \sigma_d)} H(\sigma_u - \sigma) H(\sigma - \sigma_d), \quad (84)$$

where

$$x = \frac{\sigma - \sigma_d}{\sigma_u - \sigma_d}, \quad \nu = \frac{1 - \tilde{\sigma}_0}{D_\sigma} - \tilde{\sigma}_0, \quad \mu = \left(\frac{1}{\tilde{\sigma}_0} - 1 \right) \nu, \quad \tilde{\sigma}_0 = \frac{\sigma_0 - \sigma_d}{\sigma_u - \sigma_d}, \quad (85)$$

$$\sigma_0 = \langle \sigma \rangle, \quad D_\sigma = \frac{\langle \sigma^n \rangle}{\sigma_0^2} - 1, \quad \langle \sigma^n \rangle = \int d\sigma^n \sigma F(\sigma). \quad (86)$$

Here σ_u and σ_d defined the range of values for σ . In our calculations we have chosen $\sigma_u = 1.2947\sigma_0$ and $\sigma_d = 0.85\sigma_0$.

In Figs 1-3 the liquid-gas phase diagram for polydisperse dipolar hard-sphere fluid at different degree of polydispersity $D_\sigma = 0.1, 0.2, 0.3$ is presented in T^* vs ρ^* coordinate frame. Here $T^* = k_B T / d_\mu^2(\sigma_0)$ and $\rho^* = \rho * \sigma_0^3$. We show the cloud and shadow curves and critical binodal. In addition for the reference we include Monte-Carlo predictions for the critical point and ETPT-CF predictions for the phase diagram of monodisperse version of the model [11]. One can see that upon increasing of D_σ the region of the phase instability increases with the critical point shifting to the higher temperatures and densities. For the larger values of D_σ ($D_\sigma = 0.2, 0.3$) the low density part of the cloud curve and the high density part of the shadow curve become almost vertical. For $D_\sigma = 0.3$ the cloud and shadow curves do not intersect. At $T^* = 0.1715$ the cloud curve has a cusp and shadow curve has a jump discontinuity. We believe that at this temperature there is a three-phase equilibria, when the mother phase is in equilibrium with two phases on two branches of the shadow curve, one with slightly lower density and the other with slightly higher density, respectively. The cloud and shadow curves for the whole set of values for D_σ are collected in Fig.4. In Figs 5-7 and Figs 8-10 we present the the average A -size L_A and B -size L_B of the clusters, respectively, along the cloud and shadow curves and along binodals for both polydisperse and monodisperse versions of the model. For L_K we have used expression (38), extended to account for polydispersity, i.e.

$$L_K = \int d\sigma F(\sigma) \frac{4 - [1 + \kappa_{K_1}^2(\sigma)] [1 - \kappa_{K_1}^2(\sigma)]}{[1 + \kappa_{A_1}^2(\sigma)] [1 + \kappa_{B_1}^2(\sigma)]} \left\{ \int d\sigma F(\sigma) \frac{4 - (1 - \kappa_{K_1}(\sigma)) [1 + \kappa_{K_1}^2(\sigma)]}{[1 + \kappa_{A_1}^2(\sigma)] [1 + \kappa_{B_1}^2(\sigma)]} \right\}^{-1}. \quad (87)$$

From these figures one can see that clusters of the larger sizes occurs in the liquid phase, in comparison with the gas phase. Decrease of the temperature causes increase of the cluster sizes in the liquid phase and decrease in the gas phase. With the increase of polydispersity L_K along the cloud curve do not change much. However corresponding changes along the shadow curve is more substantial, here the cluster sizes increases with polydispersity increase. In Figs 11-13 we show the ratio of A - and B -sizes L_A/L_B . As one would expect L_A is substantially larger then

L_B , thus prevailing is a chain structure of the clusters formed, with chains mutually connected via B -bonds. This difference in L_A and L_B is larger in the liquid phase and increases with polydispersity increase. Similar, as before, the temperature decrease causes L_A/L_B decrease in the gas phase and increase in the liquid phase. In Figs 14-16 and Figs 17 and 18 distribution functions of the shadow curve and on the critical binodal at different temperatures are demonstrated. According to these figures the larger size particles always fractionate to the liquid phase and smaller particles prefer to move to the gas phase. With increase of D_σ and decrease of the temperature this fractionation effects becomes more pronounced. Finally in Fig 19 we show distribution functions of the two branches of the shadow curve at $T^* = 0.1715$ and mother phase distribution function.

V. CONCLUSIONS

In this paper we propose extension of our TPT-CF approach to account for several associating sites with the possibility of each site to be multiply bonded. The theory is applied to study the liquid-gas phase behavior of polydisperse dipolar hard-sphere fluid. We present a full phase diagram, which include cloud and shadow curves, binodals and distribution functions of the coexisting phases.

VI. APPENDIX A

Orientationally averaged Mayer function $\bar{f}_{KK}(r, \sigma_1 \sigma_2)$ were fitted empirically as a sum of Yukawa-like terms

$$f_m(r, \sigma_i \sigma_j) = \sum_{n=1}^N A_n^{(m)}(\sigma_i, T) A_n^{(m)}(\sigma_j, T) \frac{e^{-z_n^{(m)}(T)(r-\sigma_{ij})}}{z_n^{(m)}(T) r}, \quad (88)$$

where $f_1 = \bar{f}_{AA}$ and $f_2 = \bar{f}_{BB}$, i, j denotes the species of the particles, $\sigma_{ij} = (\sigma_i + \sigma_j)/2$, $N = 6$, and $A_n(\sigma, T)$ are parameters that depend on particle size and temperature and were fitted by “polynomially-exponential” functions of different forms for first and second integrals.

The fitting was performed for following range of parameters: $\sigma \in [0.85, 1.2947]$ and $T \in [0.13, 0.2]$. The functional dependence of $A_n(\sigma_i, T)$ and $z_n(T)$ was chosen

differently for first and second integrals. For the **first** integral

$$\begin{aligned} A_n^{(1)}(\sigma, T) = & a_{n,1}^{(1)}(T) + a_{n,2}^{(1)}(T)x + a_{n,3}^{(1)}(T)x^2 + a_{n,4}^{(1)}(T)x^3 \\ & + a_{n,5}^{(1)}(T)e^{a_{n,6}^{(1)}(T)x + a_{n,7}^{(1)}(T)x^2 + a_{n,8}^{(1)}(T)x^6 + a_{n,9}^{(1)}(T)x^8}, \end{aligned} \quad (89)$$

where $x = (\sigma - \sigma_{min})/(\Delta\sigma)$, $\sigma_{min} = 0.85$, $\Delta\sigma = 1.2947 - \sigma_{min} = 0.4447$ and $a_{n,1}^{(1)}(T), \dots, a_{n,9}^{(1)}(T)$ are given below

$$\begin{aligned} a_{n,i}^{(1)}(T) = & e^{b_{n,i,1}^{(1)}y} \left(b_{n,i,2}^{(1)} + b_{n,i,3}^{(1)}y + b_{n,i,4}^{(1)}y^2 \right), \quad \text{for } i = 1, 2, 3, 4 \\ a_{n,i}^{(1)}(T) = & \left(b_{n,i,1}^{(1)} + b_{n,i,2}^{(1)}y + b_{n,i,3}^{(1)}y^2 + b_{n,i,4}^{(1)}y^3 \right), \quad \text{for } i = 5, 6, 7, 8, 9, \end{aligned} \quad (90)$$

where $y = \frac{T_{min}}{T}$ and $T_{min} = 0.13$.

The temperature dependence of $z_n^{(1)}(T)$ reads

$$z_n^{(1)}(T) = \left(\omega_{n,1}^{(1)} + \omega_{n,2}^{(1)}y + \omega_{n,3}^{(1)}y^2 + \omega_{n,4}^{(1)}y^3 \right). \quad (91)$$

For the **second** integral the functional dependence of $A_n^{(2)}(\sigma, T)$ and $z_n^{(2)}(T)$ is following:

$$\begin{aligned} A_n^{(2)}(\sigma) = & a_{n,1}^{(2)} + a_{n,2}^{(2)}x + a_{n,3}^{(2)}x^2 + a_{n,4}^{(2)}x^3 + a_{n,5}^{(2)}e^{a_{n,6}^{(2)}x + a_{n,7}^{(2)}x^2 + a_{n,8}^{(2)}x^6 + a_{n,9}^{(2)}x^8}, \\ z_n^{(2)}(T) = & \left(\omega_{n,1}^{(2)} + \omega_{n,2}^{(2)}y + \omega_{n,3}^{(2)}y^2 + \omega_{n,4}^{(2)}e^{\omega_{n,5}^{(2)}y} \right) \end{aligned} \quad (92)$$

where

$$a_{n,i}^{(2)} = e^{b_{n,i,1}^{(2)}y} \left(b_{n,i,2}^{(2)} + b_{n,i,3}^{(2)}y + b_{n,i,4}^{(2)}e^{b_{n,i,5}^{(2)}y} \right), \quad \text{for } i = 1, 2, 3, 4 \quad (93)$$

and

$$a_{n,i}^{(2)} = \left(b_{n,i,1}^{(2)} + b_{n,i,2}^{(2)}y + b_{n,i,3}^{(2)}y^2 + b_{n,i,4}^{(2)}e^{b_{n,i,5}^{(2)}y} \right), \quad \text{for } i = 5, 6, 7, 8, 9. \quad (94)$$

The x and y are the same as for the **first** integral.

The fitting procedure consisted of finding suitable $b_{n,i,j}$ and $\omega_{n,j}$ parameters for first and second integrals by means of differential evolution optimization algorithm [17, 18]. The objective function in both cases was a sum of square deviations of the area under $f_m(r, \sigma_i \sigma_j)$ as a function of r , and contact value $f_m(r = \sigma_{ij}, \sigma_i \sigma_j)$, from their fitting representations for different values of σ_i , σ_j and T . The deviations in the objective function were calculated for ten different temperature values (uniformly distributed from 0.13 to 0.2) and twenty different σ values (also uniformly distributed from 0.85 to 1.2947).

VII. APPENDIX B

$$m_l(\sigma) = \sigma^l, \quad (95)$$

$$m_{K,l}^{(n)}(\sigma) = \sigma^l \varphi(z_K^{(n)}, \sigma), \quad (96)$$

$$m_{K,l}^{(n0)}(\sigma) = -\frac{2\sigma^l A_K^{(n)}(\sigma)}{1 + \kappa_K^2(\sigma)} \left(1 - \frac{2\kappa_K^2(\sigma)}{1 + \kappa_K^2(\sigma)}\right), (1 - \kappa_K(\sigma)) \quad (97)$$

$$m_{K,l}^{(nm)}(\sigma) = \frac{2\sigma^l A_K^{(n)}(\sigma) A_K^{(m)}(\sigma)}{1 + \kappa_K^2(\sigma)} \left(1 - \frac{2\kappa_K^2(\sigma)}{1 + \kappa_K^2(\sigma)}\right), \quad (98)$$

$$m_\mu(\sigma) = \sum_{K=A}^B \left[1 + 2\kappa_K^2(\sigma) \frac{1 - \kappa_K(\sigma)}{1 + \kappa_K^2(\sigma)}\right] \frac{1 - \kappa_K(\sigma)}{1 + \kappa_K^2(\sigma)}, \quad (99)$$

$$\frac{\delta P_{K,1}^{(n)}}{\delta F(\sigma)} = -\frac{\rho}{2z_K^{(n)} D_K^{(n)}} \left[\pi \sigma \left(\sigma^2 + \varphi(z_K^{(n)}, \sigma) \right) + 2z_K^{(n)} P_{K,1}^{(n)} \frac{\delta D_K^{(n)}}{\delta F(\sigma)} \right], \quad (100)$$

$$\frac{\delta P_{K,2}^{(n)}}{\delta F(\sigma)} = \frac{\rho}{4z_K^{(n)} D_K^{(n)}} \left[\pi \left(\sigma^2 + 2\varphi(z_K^{(n)}, \sigma) \right) - 4z_K^{(n)} P_{K,2}^{(n)} \frac{\delta D_K^{(n)}}{\delta F(\sigma)} \right], \quad (101)$$

$$\frac{\delta P_{K,3}^{(n)}}{\delta F(\sigma)} = \frac{\rho}{(2z_K^{(n)})^2 D_K^{(n)}} \left[\pi \sigma^2 \left(\frac{2}{3} \sigma + z_K^{(n)} \varphi(z_K^{(n)}, \sigma) \right) - 2(z_K^{(n)})^2 P_{K,3}^{(n)} \frac{\delta D_K^{(n)}}{\delta F(\sigma)} \right], \quad (102)$$

$$\begin{aligned} \frac{\delta D_K^{(n)}}{\delta F(\sigma)} = & \frac{1}{6} \pi \sigma^3 \left\{ m_{K,1}^{(n)} - \frac{3}{z_K^{(n)}} \left(m_{K,0}^{(n)} + \frac{1}{2} m_2 \right) - \Delta \sigma \left(\frac{1}{6} \sigma^2 + \varphi(z_K^{(n)}, \sigma) \right) \right. \\ & - \left(\frac{1}{2} \sigma^2 + \varphi(z_K^{(n)}, \sigma) \right) \left[\frac{1}{z_K^{(n)}} \left(1 + \frac{1}{2} \pi m_3 \right) + \frac{1}{2} \pi m_{K,2}^{(n)} \right] \\ & \left. - \frac{1}{4} \pi \sigma^2 \varphi(z_K^{(n)}) \left(m_2 + 2m_{K,0}^{(n)} \right) + \frac{\pi}{2} \sigma m_{K,1}^{(n)} \varphi(z_K^{(n)}, \sigma) \right\}, \end{aligned} \quad (103)$$

$$\varphi(z, \sigma) = \frac{1}{z} (1 - z\sigma - e^{z\sigma}), \quad (104)$$

$$\begin{aligned} [M_{1,1}^{(K)}]_{nm} &= \delta_{nm} - P_{K,1}^{(m)} m_{K,1}^{(nm)} + P_{K,3}^{(m)} m_{K,0}^{(nm)} \\ [M_{1,2}^{(K)}]_{nm} &= -P_{K,1}^{(m)} m_{K,0}^{(nm)} + P_{K,2}^{(m)} m_{K,1}^{(nm)} \\ [M_{2,1}^{(K)}]_{nm} &= -P_{K,1}^{(m)} m_{K,2}^{(nm)} + P_{K,3}^{(m)} m_{K,1}^{(nm)} \\ [M_{2,2}^{(K)}]_{nm} &= \delta_{nm} - P_{K,1}^{(m)} m_{K,1}^{(nm)} + P_{K,2}^{(m)} m_{K,2}^{(nm)} \end{aligned}$$

-
- [1] Y.V.Kalyuzhnyi, G.Stell, Mol. Phys. **78**, 1247(1993).
 - [2] Y.V.Kalyuzhnyi, I.A.Protsykevitch, P.T.Cummings., Europhys. Lett. **80**, 56002(2007).
 - [3] Y.V.Kalyuzhnyi, I.A.Protsykevitch, G.Ganzenmuller, P.J.Camp, Europhys.Lett. **84**, 26001(2008).
 - [4] Y.V.Kalyuzhnyi, I.A.Protsykevitch, P.T.Cummings, J.Cond.Matt. **10**, 553(2007).
 - [5] M.S.Wertheim, J. Stat. Phys. **35**, 19;35(1984).
 - [6] M.S.Wertheim, J. Stat. Phys. **42**, 477;495(1986).
 - [7] A.Ben-Naim, J.Chem.Phys. **54**, 3682(1971).
 - [8] W.G.Chapman, G.Jackson, K.E.Gubbins, Mol.Phys. **65**, 1057(1988).
 - [9] Y.V.Kalyuzhnyi, V.Vlachy, M.F.Holovko, G.Stell, J.Chem.Phys. **102**, 5770(1995).
 - [10] T.Thusty, S.A.Safran, Science **290**, 1328(2000).
 - [11] P.J.Camp, J.C.Shelley, G.N.Patey, Phys. Rev. Lett. **84**, 115(2000).
 - [12] L.Blum, J.Hoye, J.Phys.Chem. **81**, 1311(1977).
 - [13] G.A.Mansoori, N.F.Carnahan, K.E.Startling, T.W.Leland, J.Chem.Phys. **54**, 1523(1971).
 - [14] P.Sollich, J.Phys.:Cond.Matt. **14**, R79(2002).
 - [15] L.Bellier-Castella, H.Xu, M.Baus, J.Chem.Phys. **113**, 8337(2000).
 - [16] Y.V.Kalyuzhnyi, G.Kahl, J.Chem.Phys. **119**, 7335(2003).
 - [17] R. Storn, K. Price, *Journal of global optimization*, **11**, 341 (1997).
 - [18] K.V. Price, R.M. Storn, J.A. Lampinen, *Differential evolution: a practical approach to global optimization*, Springer Verlag, 2005.




FIG. 1: Predictions of the ETPT-CF for the phase diagram of polydisperse dipolar hard-sphere mixture including cloud and shadow curves (as labeled), and critical binodal (dashed line) at $D_\sigma = 0.1$ in ρ^* vs T^* coordinate frame. Dotted line and empty square represent ETEP-CF binodal and MC [?] critical point of monodisperse dipolar hard-sphere fluid, respectively




FIG. 2: The same as in Fig.1 at $D_\sigma = 0.2$.




FIG. 3: The same as in Fig.1 at $D_\sigma = 0.3$.




FIG. 4: ETPT-CF cloud and shadow curves for polydisperse dipolar hard-sphere mixture at $D_\sigma = 0.1$ (solid lines), $D_\sigma = 0.2$ (dashed lines) and $D_\sigma = 0.3$ (dotted lines).




FIG. 5: Average A -size of the clusters along the cloud and shadow curves (as labeled), along the critical binodal (dashed line) for polydisperse dipolar hard-sphere fluid at $D_\sigma = 0.1$ and along the binodal for monodisperse version of the model (dotted lines).




FIG. 6: The same as in Fig.8 at $D_\sigma = 0.2$




FIG. 7: The same as in Fig.8 at $D_\sigma = 0.3$




FIG. 8: Average B -size of the clusters along the cloud and shadow curves (as labeled), along the critical binodal (dashed line) for polydisperse dipolar hard-sphere fluid at $D_\sigma = 0.1$ and along the binodal for monodisperse version of the model (dotted lines).




FIG. 9: The same as in Fig.11 at $D_\sigma = 0.2$.




FIG. 10: The same as in Fig.11 at $D_\sigma = 0.3$




FIG. 11: The ratio of the average A and B sizes of the clusters along the cloud and shadow curves (as labeled), along the critical binodal (dashed line) for polydisperse dipolar hard-sphere fluid at $D_\sigma = 0.1$




FIG. 12: The same as in Fig.14 at $D_\sigma = 0.2$




FIG. 13: The same as in Fig.14 at $D_\sigma = 0.3$




FIG. 14: Distribution functions of the gas (dotted lines) and liquid (dashed lines) phases along the shadow curve for $T^* = 0.16, 0.15, 0.14$ and mother phase distribution function (solid line) at $D_\sigma = 0.1$. With the temperature decrease distribution functions of the liquid phase shifts in the direction of larger σ and distribution functions of the gas phase shifts in the direction of smaller σ .




FIG. 15: The same as in Fig.17 for $T^* = 0.17, 0.16, 0.15, 0.14$ and $D_\sigma = 0.2$




FIG. 16: The same as in Fig.17 for $T^* = 0.18, 0.17, 0.16, 0.15, 0.14$ and $D_\sigma = 0.3$




FIG. 17: Distribution functions of the gas (dotted lines) and liquid (dashed lines) phases along the critical binodal for $T^* = 0.16, 0.15, 0.14$ and mother phase distribution function (solid line) at $D_\sigma = 0.1$. With the temperature decrease distribution functions of the liquid phase slightly shifts in the direction of larger σ (almost coinciding with the mother phase distribution function) and distribution functions of the gas phase shifts in the direction of smaller σ .




FIG. 18: The same as in Fig.20 at $D_\sigma = 0.2$




FIG. 19: Distribution functions on a shadow curves at $T^* = 0.1715$ for three phases at equilibria for $D_\sigma = 0.3$